

PRODUCT NO.

# 08055SUD



## STERILEX® ULTRA DISINFECTANT CLEANER SOLUTION 1

For disinfection and non-food contact sanitization for public health uses, this is part one of a two-part product and must be used with Sterilex Ultra Activator Solution.

For biocide and non-public health uses, the use of Sterilex Ultra Activator Solution may not be required.

### Powered by Patented PerQuat® Technology

#### One-Step Disinfectant | Cleaner | Sanitizer

##### EFFECTIVE AGAINST:

*Enterobacter aerogenes*  
*Staphylococcus aureus*  
*Pseudomonas aeruginosa* (*Pseudomonas*)  
*Staphylococcus aureus* (*Staph*)  
*Staphylococcus aureus* Methicillin Resistant (MRSA)  
*Salmonella enterica* (*Salmonella*)

*Escherichia coli* O157:H7  
*Listeria monocytogenes*  
*Campylobacter jejuni*  
*Cronobacter sakazakii*  
Feline calicivirus  
Norovirus  
Rotavirus

Avian influenza A Virus  
Newcastle disease virus  
Porcine respiratory & reproductive syndrome virus (PRRS)  
Mouse Parvovirus type 1 (MPV-1)  
Minute virus of Mouse (Parvovirus)

Please consult the directions for use booklet for specific instructions.

##### ACTIVE INGREDIENTS:

n-Alkyl (C <sub>14</sub> 60%, C <sub>16</sub> 30%, C <sub>12</sub> 5%, C <sub>18</sub> 5%) dimethylbenzylammonium chloride.....	3.00%
n-Alkyl (C <sub>12</sub> 68%, C <sub>14</sub> 32%) dimethylethylbenzylammonium chloride .....	3.00%
Hydrogen Peroxide .....	6.30%
Inert Ingredients .....	87.70%
<b>TOTAL .....</b>	<b>100.00%</b>

### KEEP OUT OF REACH OF CHILDREN DANGER

See side panel for additional precautionary statements.

#### FIRST AID

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control center or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**GENERAL INFORMATION:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies, call the Poison Control Center at 1.800.222.1222. For general information on this product, call 1.800.511.1659 (9 AM to 5 PM ET), or contact the National Pesticides Information Center (NPIC) at 1.800.858.7378, Monday through Friday, 8 AM to 12 PM PT, or at <http://npic.orst.edu>.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

#### PRECAUTIONARY STATEMENTS

##### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**DANGER:** Corrosive. Causes irreversible eye damage and skin burns. May be fatal if absorbed by the skin. Harmful if swallowed or inhaled. Do not get in eyes, on skin or on clothing. Avoid breathing dust, vapor or spray mist. Wear appropriate protective eyewear such as goggles, face shield, or safety glasses. Wear coveralls over long-sleeved shirt and long pants, socks, chemical resistant shoes and gloves, and when mixing and loading wear a chemical-resistant apron. Wash thoroughly with soap and water after handling and before eating, drinking, and chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

**ENVIRONMENTAL HAZARDS:** This pesticide is toxic to fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

**PHYSICAL AND CHEMICAL HAZARDS:** When exposed to fire, Sterilex Ultra Disinfectant Cleaner Solution 1 may release oxygen and oxides of carbon and/or nitrogen.

#### DIRECTIONS FOR USE (See Directions for Use)

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For disinfection and sanitization for public health uses, this is part one of a two-part product and must be used with Sterilex Ultra Activator Solution. For biocide and non-public health uses, the use of Sterilex Ultra Activator Solution may not be required. Please consult the directions for use for specific instructions.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Keep product in closed original container when not in use. Store in a cool area away from potential sources of heat, open flames, sunlight or other chemicals. Store product below 90°F.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray or mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. If rinsate cannot be used, follow pesticide disposal instructions. If not triple rinsed these containers are acute hazardous wastes and must be disposed in accordance with local, state and federal regulations. Triple rinse or pressure wash container (or equivalent) promptly after emptying.

**Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the containers on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.



**STERILEX®**  
Innovative Solutions for Microbial Control

**Manufactured by:**  
Sterilex  
111 Lake Front Drive  
Hunt Valley, MD 21030-2216

Phone (Toll-Free):  
1.800.511.1659  
Phone: 443.541.8800

EPA Reg. No.: 63761-8

Revision: 11112022

**NET WEIGHT:  
460 LB**

PRODUCT NO.

# 08055SUD

## STERILEX® ULTRA DISINFECTANT CLEANER SOLUTION 1

